

## **QCE Chemistry Units 3&4**

### **Paper 1**

Student's Name: \_\_\_\_\_

Teacher's Name: \_\_\_\_\_

#### **Time allowed**

- Perusal time – 10 minutes
- Working time – 90 minutes

#### **General instructions**

- Answer all questions in this question and response booklet.
- QCAA-approved calculator permitted.
- Formula and data booklet provided.
- Planning paper will not be marked.

#### **Section 1 (25 marks)**

- 25 multiple choice questions

#### **Section 2 (35 marks)**

- 8 short response questions

Students are advised that this is a trial examination only and cannot in any way guarantee the content or the format of the 2022 QCE Chemistry Units 3&4 Written Examination.

Neap<sup>®</sup> Education (Neap) Trial Exams are licensed to be photocopied or placed on the school intranet and used only within the confines of the school purchasing them, for the purpose of examining that school's students only. They may not be otherwise reproduced or distributed. The copyright of Neap Trial Exams remains with Neap. No Neap Trial Exam or any part thereof is to be issued or passed on by any person to any party inclusive of other schools, non-practising teachers, coaching colleges, tutors, parents, students, publishing agencies or websites without the express written consent of Neap.

**SECTION 1****Instructions**

- Choose the best answer for Questions 1–25.
- This section has 25 questions and is worth 25 marks.
- Use a 2B pencil to fill in the A, B, C or D answer bubble completely.
- If you change your mind or make a mistake, use an eraser to remove your response and fill in the new answer bubble completely.

	A	B	C	D
Example:	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>


	A	B	C	D
<b>1.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>2.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>3.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>4.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>5.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>6.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>7.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>8.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>9.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>10.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>11.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>12.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>13.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>14.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>15.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>16.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>17.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>18.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>19.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>20.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>21.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>22.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>23.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>24.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<b>25.</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

**SECTION 2****Instructions**

- Write using black or blue pen.
  - If you need more space for a response, use the additional pages at the back of this booklet.
    - On the additional pages, write the question number you are responding to.
    - Cancel any incorrect response by ruling a single diagonal line through your work.
    - Write the page number of your alternative/additional response, i.e. See page ...
    - If you do not do this, your original response will be marked.
  - This section has eight questions and is worth 35 marks.
- 

**QUESTION 26 (2 marks)**

Sulfur dioxide and oxygen are mixed to produce sulfur trioxide according to the following equation.



A catalyst, vanadium pentoxide, is then added to the mixture.

- a) Predict the effect that the catalyst will have on the equilibrium yield. *[1 mark]*

---

---


- b) Predict the effect that the catalyst will have on the rate of reaction. *[1 mark]*

---

---

**QUESTION 27 (3 marks)**

5 mL of a solution containing  $\text{Fe}^{3+}$  ions is added to a 10 mL of a solution containing  $\text{SCN}^-$  ions in a large test tube at room temperature. The reaction occurs according to the following equation.



The resulting solution is pale red.

- a) Another 5 mL of the solution containing  $\text{Fe}^{3+}$  ions is added to the test tube, and the colour of the resulting solution becomes darker.

Explain why the colour change occurred.

[1 mark]

---

---

- b) The test tube from 27a) is placed in a water bath at approximately  $90^\circ\text{C}$ . The colour of the solution in the test tube becomes lighter, changing to yellow.

Deduce whether the reaction is endothermic or exothermic.

[2 marks]

---


---

---


---

**QUESTION 28 (5 marks)**

Hydrazine,  $\text{N}_2\text{H}_4$ , is a chemical that is used mainly as a foaming agent in the preparation of polymer foams and as a long-term storable propellant for in-space propulsion of spacecraft. It can be produced through several reactions; two of these reactions are shown.

**Reaction 1**

ammonia + sodium hypochlorite  $\rightarrow$  hydrazine + sodium chloride + water

**Reaction 2**

chloramine + excess ammonia  $\rightarrow$  hydrazine + ammonium chloride

Determine which reaction has better atom economy and is thus more efficient. Show your working.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

---

**QUESTION 29 (4 marks)**

The diagram shows two amino acids that have bonded to form a dipeptide.



- a) Identify the two amino acids. [2 marks]


---



---

- b) On the diagram above, circle the peptide bond that joins the amino acid molecules. [1 mark]

- c) An amino acid is shown.



- Determine why this amino acid can exist as an anion with a charge of 2-. [1 mark]


---



---

**QUESTION 30 (7 marks)**

The flow chart shows a series of reactions involving four compounds: A, B, C and D.



a) Sketch the structural formulas of compounds A, C and D.

[3 marks]

Compound	Structural formula
A	
C	
D	

- b) The type of reaction that produces compound B is different to the type of reaction that produces compound D.

Identify and describe the two types of reaction.

[2 marks]

Reaction that produces compound B \_\_\_\_\_

\_\_\_\_\_

Reaction that produces compound D \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

- c) Compound B has a boiling point of 97°C and compound C has a boiling point of 141.2°C.

Explain the difference in boiling points.

[2 marks]

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_


\_\_\_\_\_

\_\_\_\_\_



**QUESTION 31 (4 marks)**

The flow chart shows the steps in a particular industrial process. The conditions for reaction 2, such as temperature and pressure, can be altered.



- a) If the temperature of the vessel for reaction 2 is lowered, explain the effect this will have on the yield of the reaction and the value of the equilibrium constant ( $K_c$ ). [2 marks]

---



---



---



---

- b) If the pressure of the vessel for reaction 2 is increased to 5 atm by pumping more C(g) and B(g) into the vessel at constant temperature, infer the effect this will have on the yield of the reaction and the value of  $K_c$ . [2 marks]

---



---





---



---

**QUESTION 32 (6 marks)**

The structural formulas of three acids that are commonly found in fruit and associated with winemaking are shown.



The table shows the dissociation constants ( $K_a$ ) and  $\text{p}K_a$  values of the acids.

Acid	Formula	$K_a$	$\text{p}K_a$ value(s)
ethanoic acid	$\text{CH}_3\text{COOH}$	$1.75 \times 10^{-5}$	4.75
tartaric acid	$\text{C}_4\text{H}_6\text{O}_6$	$1.0 \times 10^{-3}$	2.95, 4.25
malic acid	$\text{C}_4\text{H}_6\text{O}_5$	$4.0 \times 10^{-4}$	3.4, 5.2

- a) Using the first  $\text{p}K_a$  value of each acid, determine which acid is the strongest. [1 mark]

---



---

- b) Propose why malic acid has two  $\text{p}K_a$  values. [1 mark]

---



---



---

- c) Calculate the pH of a 0.50 M solution of ethanoic acid. Show your working. In your response, include a balanced chemical equation for the ionisation of ethanoic acid and state any assumptions made.

[4 marks]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

pH = \_\_\_\_\_ (to two decimal places)

**QUESTION 33 (4 marks)**

A student titrated a solution of a weak acid of unknown concentration with a 1.0 M solution of standard sodium hydroxide, NaOH. The pH of the acid solution was monitored as the titration proceeded and is shown in the graph.



- a) What was the initial pH of the acid solution? *[1 mark]*
- 
- b) On the graph, label the equivalence point, half-equivalence point and buffer zone. *[3 marks]*

**END OF PAPER**













Trial Examination 2022

---

**Formula and Data Booklet**

# **QCE Chemistry Units 3&4**

---

Neap<sup>®</sup> Education (Neap) Trial Exams are licensed to be photocopied or placed on the school intranet and used only within the confines of the school purchasing them, for the purpose of examining that school's students only. They may not be otherwise reproduced or distributed. The copyright of Neap Trial Exams remains with Neap. No Neap Trial Exam or any part thereof is to be issued or passed on by any person to any party inclusive of other schools, non-practising teachers, coaching colleges, tutors, parents, students, publishing agencies or websites without the express written consent of Neap.

**FORMULAS****Processing of data**

$$\text{Absolute uncertainty of the mean } \Delta\bar{x} = \pm \frac{(x_{\max} - x_{\min})}{2}$$

$$\text{Percentage uncertainty (\%)} = \frac{\text{absolute uncertainty}}{\text{measurement}} \times \frac{100}{1}$$

$$\text{Percentage error (\%)} = \left| \frac{\text{measured value} - \text{true value}}{\text{true value}} \right| \times 100$$

**Chemical reactions – reactants, products and energy change**

$$\Delta H = H_{(\text{products})} - H_{(\text{reactants})}$$

$$\Delta H = \Sigma(\text{bonds broken}) - \Sigma(\text{bonds formed})$$

$$Q = mc\Delta T$$

$$\text{Percentage yield (\%)} = \frac{\text{experimental yield}}{\text{theoretical yield}} \times \frac{100}{1}$$

**Aqueous solutions and acidity**

$$\text{Molarity} = \frac{\text{moles of solute (n)}}{\text{volume of solution (V)}}$$

**Chemical equilibrium systems**

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b} \text{ for the reaction: } aA + bB \rightleftharpoons cC + dD$$

$$K_w = [H^+][OH^-]$$

$$\text{pH} = -\log_{10} [H^+]$$

$$\text{pOH} = -\log_{10} [OH^-]$$

$$K_w = K_a \times K_b$$

$$K_a = \frac{[H_3O^+][A^-]}{[HA]}$$

$$K_b = \frac{[BH^+][OH^-]}{[B]}$$

**PHYSICAL CONSTANTS AND UNIT CONVERSIONS**

Physical constants and unit conversions	
Absolute zero	$0 \text{ K} = -273^\circ\text{C}$
Atomic mass unit	$1 \text{ amu} = 1.66 \times 10^{-27} \text{ kg}$
Avogadro's constant	$N_{\text{A}} = 6.02 \times 10^{23} \text{ mol}^{-1}$
Ideal gas constant	$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
Ionic product constant for water (at 298 K)	$K_{\text{w}} = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$
Molar volume of an ideal gas (at STP)	$2.27 \times 10^{-2} \text{ m}^3 \text{ mol}^{-1} = 22.7 \text{ dm}^3 \text{ mol}^{-1}$
Specific heat capacity of water (at 298 K)	$c_{\text{w}} = 4.18 \text{ J g}^{-1} \text{ K}^{-1}$
Standard temperature and pressure (STP)	273 K and 100 kPa
Volume and capacity conversions	$1 \text{ dm}^3 = 1 \times 10^{-3} \text{ m}^3 = 1 \times 10^3 \text{ cm}^3 = 1 \text{ L}$

**LIST OF ELEMENTS**

Name	Atomic no.	Symbol
Hydrogen	1	H
Helium	2	He
Lithium	3	Li
Beryllium	4	Be
Boron	5	B
Carbon	6	C
Nitrogen	7	N
Oxygen	8	O
Fluorine	9	F
Neon	10	Ne
Sodium	11	Na
Magnesium	12	Mg
Aluminium	13	Al
Silicon	14	Si
Phosphorus	15	P
Sulfur	16	S
Chlorine	17	Cl
Argon	18	Ar
Potassium	19	K
Calcium	20	Ca
Scandium	21	Sc
Titanium	22	Ti
Vanadium	23	V
Chromium	24	Cr
Manganese	25	Mn
Iron	26	Fe
Cobalt	27	Co
Nickel	28	Ni
Copper	29	Cu
Zinc	30	Zn
Gallium	31	Ga
Germanium	32	Ge
Arsenic	33	As
Selenium	34	Se
Bromine	35	Br

Name	Atomic no.	Symbol
Krypton	36	Kr
Rubidium	37	Rb
Strontium	38	Sr
Yttrium	39	Y
Zirconium	40	Zr
Niobium	41	Nb
Molybdenum	42	Mo
Technetium	43	Tc
Ruthenium	44	Ru
Rhodium	45	Rh
Palladium	46	Pd
Silver	47	Ag
Cadmium	48	Cd
Indium	49	In
Tin	50	Sn
Antimony	51	Sb
Tellurium	52	Te
Iodine	53	I
Xenon	54	Xe
Cesium	55	Cs
Barium	56	Ba
Lanthanum	57	La
Cerium	58	Ce
Praseodymium	59	Pr
Neodymium	60	Nd
Promethium	61	Pm
Samarium	62	Sm
Europium	63	Eu
Gadolinium	64	Gd
Terbium	65	Tb
Dysprosium	66	Dy
Holmium	67	Ho
Erbium	68	Er
Thulium	69	Tm
Ytterbium	70	Yb

**LIST OF ELEMENTS (CONTINUED)**

Name	Atomic no.	Symbol
Lutetium	71	Lu
Hafnium	72	Hf
Tantalum	73	Ta
Tungsten	74	W
Rhenium	75	Re
Osmium	76	Os
Iridium	77	Ir
Platinum	78	Pt
Gold	79	Au
Mercury	80	Hg
Thallium	81	Tl
Lead	82	Pb
Bismuth	83	Bi
Polonium	84	Po
Astatine	85	At
Radon	86	Rn
Francium	87	Fr
Radium	88	Ra
Actinium	89	Ac
Thorium	90	Th
Protactinium	91	Pa
Uranium	92	U
Neptunium	93	Np
Plutonium	94	Pu

Name	Atomic no.	Symbol
Americium	95	Am
Curium	96	Cm
Berkelium	97	Bk
Californium	98	Cf
Einsteinium	99	Es
Fermium	100	Fm
Mendelevium	101	Md
Nobelium	102	No
Lawrencium	103	Lr
Rutherfordium	104	Rf
Dubnium	105	Db
Seaborgium	106	Sg
Bohrium	107	Bh
Hassium	108	Hs
Meitnerium	109	Mt
Darmstadtium	110	Ds
Roentgenium	111	Rg
Copernicium	112	Cn
Nihonium	113	Nh
Flerovium	114	Fl
Moscovium	115	Mc
Livermorium	116	Lv
Tennessine	117	Ts
Oganesson	118	Og

**PERIODIC TABLE OF THE ELEMENTS**

		KEY																
		1 atomic number																
		H symbol																
		1.01 relative atomic mass*																
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
<b>H</b> 1.01	<b>He</b> 4.00	<b>Li</b> 6.94	<b>Be</b> 9.01	<b>B</b> 10.81	<b>C</b> 12.01	<b>N</b> 14.01	<b>O</b> 16.00	<b>F</b> 19.00	<b>Ne</b> 20.18	<b>Na</b> 22.99	<b>Mg</b> 24.31	<b>Al</b> 26.98	<b>Si</b> 28.09	<b>P</b> 30.97	<b>S</b> 32.06	<b>Cl</b> 35.45	<b>Ar</b> 39.95	
<b>K</b> 39.10	<b>Ca</b> 40.08	<b>Sc</b> 44.96	<b>Ti</b> 47.87	<b>V</b> 50.94	<b>Cr</b> 52.00	<b>Mn</b> 54.94	<b>Fe</b> 55.85	<b>Co</b> 58.93	<b>Ni</b> 58.69	<b>Cu</b> 63.55	<b>Zn</b> 65.38	<b>Ga</b> 69.72	<b>Ge</b> 72.63	<b>As</b> 74.92	<b>Se</b> 78.97	<b>Br</b> 79.90	<b>Kr</b> 83.80	
<b>Rb</b> 85.47	<b>Sr</b> 87.62	<b>Y</b> 88.91	<b>Zr</b> 91.22	<b>Nb</b> 92.91	<b>Mo</b> 95.95	<b>Tc</b> (98.91)	<b>Ru</b> 101.07	<b>Rh</b> 102.91	<b>Pd</b> 106.42	<b>Ag</b> 107.87	<b>Cd</b> 112.41	<b>In</b> 114.82	<b>Sn</b> 118.71	<b>Sb</b> 121.76	<b>Te</b> 127.60	<b>I</b> 126.90	<b>Xe</b> 131.29	
<b>Cs</b> 132.91	<b>Ba</b> 137.33	Lanthanoids	<b>Hf</b> 178.49	<b>Ta</b> 180.95	<b>W</b> 183.84	<b>Re</b> 186.21	<b>Os</b> 190.23	<b>Ir</b> 192.22	<b>Pt</b> 195.08	<b>Au</b> 196.97	<b>Hg</b> 200.59	<b>Tl</b> 204.38	<b>Pb</b> 207.2	<b>Bi</b> 208.98	<b>Po</b> (210.0)	<b>At</b> (210.0)	<b>Rn</b> (222.0)	
<b>Fr</b> (223.0)	<b>Ra</b> (226.1)	Actinoids	<b>Rf</b> (261.1)	<b>Db</b> (262.1)	<b>Sg</b> (263.1)	<b>Bh</b> (264.1)	<b>Hs</b> (265.1)	<b>Mt</b> (268)	<b>Ds</b> (281)	<b>Rg</b> (272)	<b>Cn</b> (285)	<b>Nh</b> (284)	<b>Fl</b> (289)	<b>Mc</b> (288)	<b>Lv</b> (293)	<b>Ts</b> (294)	<b>Og</b> (294)	
			Lanthanoids															
			57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	
			<b>La</b> 138.91	<b>Ce</b> 140.12	<b>Pr</b> 140.91	<b>Nd</b> 144.24	<b>Pm</b> (146.9)	<b>Sm</b> 150.36	<b>Eu</b> 151.96	<b>Gd</b> 157.25	<b>Tb</b> 158.93	<b>Dy</b> 162.50	<b>Ho</b> 164.93	<b>Er</b> 167.26	<b>Tm</b> 168.93	<b>Yb</b> 173.05	<b>Lu</b> 174.97	
			Actinoids															
			89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	
			<b>Ac</b> (227.0)	<b>Th</b> 232.0	<b>Pa</b> 231.0	<b>U</b> 238.0	<b>Np</b> (237.0)	<b>Pu</b> (239.1)	<b>Am</b> (241.1)	<b>Cm</b> (244.1)	<b>Bk</b> (249.1)	<b>Cf</b> (252.1)	<b>Es</b> (252.1)	<b>Fm</b> (252.1)	<b>Md</b> (258.1)	<b>No</b> (259.1)	<b>Lr</b> (262.1)	

Groups are numbered according to IUPAC convention 1–18.  
\*Values in brackets are for the isotope with the longest half-life.

**ATOMIC AND IONIC RADII OF SELECTED ELEMENTS**

		KEY																																																																																	
		atomic number																																																																																	
		symbol																																																																																	
		atomic radius ( $10^{-12}$ m)																																																																																	
		charge of ion																																																																																	
		ionic radius ( $10^{-12}$ m)																																																																																	
		atomic number																																																																																	
		symbol																																																																																	
		atomic radius ( $10^{-12}$ m)																																																																																	
		charge of ion																																																																																	
1	<b>H</b> 1 32 208 (1-)	2	<b>Be</b> 4 99 45 (2+)	3	<b>Li</b> 3 130 76 (1+)	4	<b>Na</b> 11 160 102 (1+)	5	<b>Mg</b> 12 140 72 (2+)	6	<b>K</b> 19 200 138 (1+)	7	<b>Ca</b> 20 174 100 (2+)	8	<b>Rb</b> 37 215 152 (1+)	9	<b>Sr</b> 38 190 118 (2+)	10	<b>Ba</b> 56 206 135 (2+)	11	<b>Sc</b> 21 159 75 (3+)	12	<b>Ti</b> 22 148 86 (2+)	13	<b>V</b> 23 144 79 (2+)	14	<b>Cr</b> 24 130 62 (3+)	15	<b>Mn</b> 25 129 83 (2+)	16	<b>Fe</b> 26 124 78 (2+)	17	<b>Co</b> 27 118 74 (2+)	18	<b>Ni</b> 28 117 69 (2+)	19	<b>Cu</b> 29 122 77 (1+)	20	<b>Zn</b> 30 120 74 (2+)	21	<b>Ga</b> 31 123 62 (3+)	22	<b>Ge</b> 32 120 53 (4+)	23	<b>As</b> 33 120 58 (3+)	24	<b>Se</b> 34 118 188 (2-)	25	<b>Br</b> 35 117 196 (1-)	26	<b>Kr</b> 36 116	27	<b>Y</b> 39 176 90 (3+)	28	<b>Zr</b> 40 164 72 (4+)	29	<b>Nb</b> 41 156 64 (5+)	30	<b>Mo</b> 42 148 65 (4+)	31	<b>Tc</b> 43 138 65 (4+)	32	<b>Ru</b> 44 136 62 (4+)	33	<b>Rh</b> 45 134 67 (3+)	34	<b>Pd</b> 46 130 86 (2+)	35	<b>Ag</b> 47 136 115 (1+)	36	<b>Cd</b> 48 140 95 (2+)	37	<b>In</b> 49 142 80 (3+)	38	<b>Sn</b> 50 140 69 (4+)	39	<b>Sb</b> 51 140 76 (3+)	40	<b>Te</b> 52 137 221 (2-)	41	<b>I</b> 53 136 220 (1-)	42	<b>Xe</b> 54 136

Groups are numbered according to IUPAC convention 1–18.

**ELECTRONEGATIVITIES AND FIRST IONISATION ENERGIES OF SELECTED ELEMENTS**

													18
													<b>2</b> <b>He</b> 2379
													17
													<b>9</b> <b>F</b> 4.0 1687
													<b>8</b> <b>O</b> 3.4 1320
													15
													<b>7</b> <b>N</b> 3.0 1407
													<b>6</b> <b>C</b> 2.6 1093
													14
													<b>5</b> <b>B</b> 2.0 807
													<b>13</b> <b>Al</b> 1.6 584
													<b>14</b> <b>Si</b> 1.9 793
													<b>15</b> <b>P</b> 2.2 1018
													<b>16</b> <b>S</b> 2.6 1006
													<b>17</b> <b>Cl</b> 3.2 1257
													<b>18</b> <b>Ar</b> 1527
													12
													<b>30</b> <b>Zn</b> 1.7 913
													<b>29</b> <b>Cu</b> 1.9 752
													<b>28</b> <b>Ni</b> 1.9 743
													<b>27</b> <b>Co</b> 1.9 765
													<b>26</b> <b>Fe</b> 1.8 766
													<b>25</b> <b>Mn</b> 1.6 724
													<b>24</b> <b>Cr</b> 1.7 659
													<b>23</b> <b>V</b> 1.6 656
													<b>22</b> <b>Ti</b> 1.5 664
													<b>21</b> <b>Sc</b> 1.4 637
													<b>48</b> <b>Cd</b> 1.7 874
													<b>47</b> <b>Ag</b> 1.9 737
													<b>46</b> <b>Pd</b> 2.2 811
													<b>45</b> <b>Rh</b> 2.3 726
													<b>44</b> <b>Ru</b> 2.2 717
													<b>43</b> <b>Tc</b> 1.9 708
													<b>42</b> <b>Mo</b> 2.2 691
													<b>41</b> <b>Nb</b> 1.6 670
													<b>40</b> <b>Zr</b> 1.3 666
													<b>39</b> <b>Y</b> 1.2 606
													<b>56</b> <b>Ba</b> 0.9 509
													<b>55</b> <b>Cs</b> 0.8 382
													<b>54</b> <b>Xe</b> 2.6 1177
													<b>53</b> <b>I</b> 2.7 1015
													<b>52</b> <b>Te</b> 2.1 876
													<b>51</b> <b>Sb</b> 2.1 840
													<b>50</b> <b>Sn</b> 2.0 715
													<b>49</b> <b>In</b> 1.8 565
													<b>32</b> <b>Ge</b> 2.0 768
													<b>31</b> <b>Ga</b> 1.8 585
													<b>34</b> <b>Se</b> 2.6 947
													<b>33</b> <b>As</b> 2.2 953
													<b>36</b> <b>Kr</b> 2.9 1357
													<b>35</b> <b>Br</b> 3.0 1146
													<b>10</b> <b>Ne</b> 2087
													<b>11</b> <b>Na</b> 0.9 502
													<b>12</b> <b>Mg</b> 1.3 744
													<b>19</b> <b>K</b> 0.8 425
													<b>20</b> <b>Ca</b> 1.0 596
													<b>37</b> <b>Rb</b> 0.8 409
													<b>38</b> <b>Sr</b> 1.0 556
													<b>4</b> <b>Be</b> 1.6 906
													<b>3</b> <b>Li</b> 1.0 526
													<b>1</b> <b>H</b> 2.2 1318

KEY

<b>1</b>	atomic number
<b>H</b>	symbol
2.2	electronegativity
1318	first ionisation enthalpies (kJ mol <sup>-1</sup> )

Groups are numbered according to IUPAC convention 1–18.



**SOLUBILITY OF SELECTED COMPOUNDS AT 298 K**

	bromide	carbonate	chloride	hydroxide	iodide	nitrate	oxide	phosphate	sulfate
<b>aluminium</b>	s	–	s	i	s	s	i	i	s
<b>ammonium</b>	s	s	s	s	s	s	–	s	s
<b>barium</b>	s	i	s	s	s	s	s	i	i
<b>calcium</b>	s	i	s	p	s	s	p	i	p
<b>cobalt(II)</b>	s	i	s	i	s	s	i	i	s
<b>copper(II)</b>	s	–	s	i	i	s	i	i	s
<b>iron(II)</b>	s	i	s	i	s	s	i	i	s
<b>iron(III)</b>	s	–	s	i	s	s	i	i	s
<b>lead(II)</b>	p	i	s	i	i	s	i	i	i
<b>lithium</b>	s	s	s	s	s	s	s	–	s
<b>magnesium</b>	s	i	s	i	s	s	i	p	s
<b>manganese(II)</b>	s	i	s	i	s	s	i	p	s
<b>potassium</b>	s	s	s	s	s	s	s	s	s
<b>silver</b>	i	i	i	i	i	s	i	i	p
<b>sodium</b>	s	s	s	s	s	s	s	s	s
<b>zinc</b>	s	i	s	i	s	s	i	i	s

**Key**

Abbreviation	Explanation
s	soluble in water (solubility greater than 10 g L <sup>-1</sup> )
p	partially soluble in water (solubility between 1 and 10 g L <sup>-1</sup> )
i	insoluble in water (solubility less than 1 g L <sup>-1</sup> )
–	no data


**AVERAGE BOND ENTHALPIES AT 298 K****Single bonds**

	$\Delta H$ (kJ mol <sup>-1</sup> )								
	H	C	N	O	F	S	Cl	Br	I
H	436								
C	414	346							
N	391	286	158						
O	463	358	214	144					
F	567	492	278	191	159				
S	364	289			327	266			
Cl	431	324	192	206	255	271	242		
Br	366	285		201	249	218	219	193	
I	298	228		201	280		211	178	151

**Multiple bonds**

Bond	$\Delta H$ (kJ mol <sup>-1</sup> )
C=C	614
C≡C	839
C=N	615
C≡N	890
C=O	804
N=N	470
N≡N	945
O=O	498



**REACTIVITY SERIES OF METALS**

Element	Reactivity
K	<p>most reactive</p>  <p>least reactive</p>
Na	
Li	
Ba	
Sr	
Ca	
Mg	
Al	
C*	
Mn	
Zn	
Cr	
Fe	
Cd	
Co	
Ni	
Sn	
Pb	
H <sub>2</sub> *	
Sb	
Bi	
Cu	
Hg	
Ag	
Au	
Pt	

\* Carbon (C) and hydrogen gas (H<sub>2</sub>) added for comparison

**STANDARD ELECTRODE POTENTIALS AT 298 K**

Oxidised species $\rightleftharpoons$ Reduced species	$E^\circ$ (V)
$\text{Li}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Li}(\text{s})$	-3.04
$\text{K}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{K}(\text{s})$	-2.94
$\text{Ba}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ba}(\text{s})$	-2.91
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ca}(\text{s})$	-2.87
$\text{Na}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Na}(\text{s})$	-2.71
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mg}(\text{s})$	-2.36
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Al}(\text{s})$	-1.68
$\text{Mn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mn}(\text{s})$	-1.18
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Zn}(\text{s})$	-0.76
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Fe}(\text{s})$	-0.44
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$	-0.24
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}(\text{s})$	-0.14
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb}(\text{s})$	-0.13
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0.00
$\text{Cu}^{2+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Cu}^+(\text{aq})$	+0.16
$\text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{SO}_2(\text{aq}) + 2\text{H}_2\text{O}(\text{l})$	+0.16
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.34
$\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightleftharpoons 4\text{OH}^-(\text{aq})$	+0.40
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.52
$\text{I}_2(\text{s}) + 2\text{e}^- \rightleftharpoons 2\text{I}^-(\text{aq})$	+0.54
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s})$	+0.80
$\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-(\text{aq})$	+1.08
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$	+1.36
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+(\text{aq}) + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$	+1.36
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$	+1.51
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-(\text{aq})$	+2.89

**GLUCOSE AND FRUCTOSE: STRAIGHT CHAIN AND  $\alpha$ -RING FORMS**


## COMMON AMINO ACIDS

Common name (symbol)	Structural formula	pH of isoelectric point	Common name (symbol)	Structural formula	pH of isoelectric point
Alanine (Ala)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_3 \end{array}$	6.1	Arginine (Arg)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_2 \\   \\ \text{CH}_2 \\   \\ \text{CH}_2 \\   \\ \text{NH} \\   \\ \text{C}=\text{NH} \\   \\ \text{NH}_2 \end{array}$	10.7
Asparagine (Asn)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_2 \\   \\ \text{C}=\text{O} \\   \\ \text{NH}_2 \end{array}$	5.4	Aspartic acid (Asp)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_2 \\   \\ \text{C}=\text{O} \\   \\ \text{OH} \end{array}$	3.0
Cysteine (Cys)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_2 \\   \\ \text{SH} \end{array}$	5.1	Glutamic acid (Glu)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_2 \\   \\ \text{CH}_2 \\   \\ \text{C}=\text{O} \\   \\ \text{OH} \end{array}$	3.2
Glutamine (Gln)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{CH}_2 \\   \\ \text{CH}_2 \\   \\ \text{C}=\text{O} \\   \\ \text{NH}_2 \end{array}$	5.7	Glycine (Gly)	$\begin{array}{c} \text{H} \quad \text{O} \\   \quad    \\ \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\   \\ \text{H} \end{array}$	6.1

**COMMON AMINO ACIDS (continued)**

Common name (symbol)	Structural formula	pH of isoelectric point	Common name (symbol)	Structural formula	pH of isoelectric point
Histidine (His)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{N} \quad \text{NH} \\  \diagup \quad \diagdown \\  \text{C} \\  \diagdown \quad \diagup \\  \text{N}  \end{array}  $	7.6	Isoleucine (Ile)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CHCH}_3 \\    \\  \text{CH}_2 \\    \\  \text{CH}_3  \end{array}  $	6.0
Leucine (Leu)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{CHCH}_3 \\    \\  \text{CH}_3  \end{array}  $	6.0	Lysine (Lys)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{NH}_2  \end{array}  $	9.7
Methionine (Met)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{S} \\    \\  \text{CH}_3  \end{array}  $	5.7	Phenylalanine (Phe)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{C}_6\text{H}_5  \end{array}  $	5.7
Proline (Pro)	$  \begin{array}{c}  \text{O} \\     \\  \text{C}-\text{OH} \\    \\  \text{HN} \\  \diagup \quad \diagdown \\  \text{C} \\  \diagdown \quad \diagup \\  \text{C} \\  \diagup \quad \diagdown \\  \text{C} \\  \diagdown \quad \diagup \\  \text{C} \\  \diagup \quad \diagdown \\  \text{HN}  \end{array}  $	6.3	Serine (Ser)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{OH}  \end{array}  $	5.7

**COMMON AMINO ACIDS (continued)**

Common name (symbol)	Structural formula	pH of isoelectric point	Common name (symbol)	Structural formula	pH of isoelectric point
Threonine (Thr)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CHOH} \\    \\  \text{CH}_3  \end{array}  $	5.6	Tryptophan (Trp)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{HN} \text{---} \text{Indole ring}  \end{array}  $	5.9
Tyrosine (Tyr)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{C}_6\text{H}_4 \\    \\  \text{OH}  \end{array}  $	5.7	Valine (Val)	$  \begin{array}{c}  \text{H} \quad \text{O} \\    \quad    \\  \text{H}_2\text{N}-\text{C}-\text{C}-\text{OH} \\    \\  \text{CHCH}_3 \\    \\  \text{CH}_3  \end{array}  $	6.0



**ACID-BASE INDICATORS**

Name	pKa	pH range of colour change	Colour change (acidic to basic)
Methyl orange	3.7	3.1–4.4	red to yellow
Bromophenol blue	4.2	3.0–4.6	yellow to blue
Bromocresol green	4.7	3.8–5.4	yellow to blue
Methyl red	5.1	4.4–6.2	pink to yellow
Bromothymol blue	7.0	6.0–7.6	yellow to blue
Phenol red	7.9	6.8–8.4	yellow to red
Phenolphthalein	9.6	8.3–10.0	colourless to pink

**INFRARED DATA**

The table below shows the characteristic range of infrared absorption due to stretching in organic molecules.

Bond	Organic molecules	Wavelength ( $\text{cm}^{-1}$ )
C–I	iodoalkanes	490–620
C–Br	bromoalkanes	500–600
C–Cl	chloroalkanes	600–800
C–F	fluoroalkanes	1000–1400
C–O	alcohol, ester	1050–1410
C=C	alkenes	1620–1680
C=O	aldehydes, carboxylic acid, ester, ketones	1700–1750
C≡C	alkynes	2100–2260
O–H	carboxylic acids (hydrogen-bonded)	2500–3000
C–H	alkanes, alkenes, alkynes, aldehydes, amides	2720–3100
O–H	alcohol (hydrogen-bonded)	3200–3600
N–H	amines	3300–3500

**FORMULAS AND CHARGES FOR COMMON POLYATOMIC IONS**

Anions		Cations	
acetate (ethanoate)	$\text{CH}_3\text{COO}^-$ or $\text{C}_2\text{H}_3\text{O}_2^-$	ammonium	$\text{NH}_4^+$
carbonate	$\text{CO}_3^{2-}$	hydronium	$\text{H}_3\text{O}^+$
chlorate	$\text{ClO}_3^-$		
chlorite	$\text{ClO}_2^-$		
chromate	$\text{CrO}_4^{2-}$		
citrate	$\text{C}_6\text{H}_5\text{O}_7^{3-}$		
cyanide	$\text{CN}^-$		
dichromate	$\text{Cr}_2\text{O}_7^{2-}$		
dihydrogen phosphate	$\text{H}_2\text{PO}_4^-$		
hypochlorite	$\text{ClO}^-$		
hydrogen carbonate	$\text{HCO}_3^-$		
hydrogen sulfate	$\text{HSO}_4^-$		
hydrogen phosphate	$\text{HPO}_4^{2-}$		
hydroxide	$\text{OH}^-$		
nitrate	$\text{NO}_3^-$		
nitrite	$\text{NO}_2^-$		
perchlorate	$\text{ClO}_4^-$		
permanganate	$\text{MnO}_4^-$		
peroxide	$\text{O}_2^{2-}$		
phosphate	$\text{PO}_4^{3-}$		
sulfate	$\text{SO}_4^{2-}$		
sulfite	$\text{SO}_3^{2-}$		
thiosulfate	$\text{S}_2\text{O}_3^{2-}$		

## REFERENCES

Aylward, G and Findlay, T 2008, *SI Chemical Data*, 5th ed, John Wiley & Sons, Brisbane.

Haynes, WM (ed) 2016, *CRC Handbook of Chemistry and Physics*, 97th ed, CRC Press, Boca Raton, US.