

---

**Question 1**

The function  $f(x) = \log_e(x - 2)$  is defined for

$$x - 2 > 0$$

$$x > 2$$

So  $d_f = (2, \infty)$

**(1 mark)**

**Question 2**

a.  $f(g(x))$  exists iff  $r_g \subseteq d_f$ .

Now,  $r_g = R$  and  $d_f = R \setminus \{0\}$ .

Since  $R \not\subseteq R \setminus \{0\}$ ,

$$r_g \not\subseteq d_f$$

so  $f(g(x))$  does not exist.

**(1 mark)**

b. i.

$$\begin{aligned} g(f(x)) &= g\left(\frac{1}{2x}\right) \\ &= \frac{1}{2x} + 1 \end{aligned}$$

**(1 mark)**

ii.

$$\begin{aligned} d_{g(f(x))} &= d_f \\ &= R \setminus \{0\} \end{aligned}$$

**(1 mark)**

**Question 3**

a.  $f : (1, \infty) \rightarrow \mathbb{R}, f(x) = \frac{3}{x-1} + 2$

Let  $y = \frac{3}{x-1} + 2$

Swap  $x$  and  $y$

$$x = \frac{3}{y-1} + 2$$

**(1 mark)**

$$x - 2 = \frac{3}{y-1}$$

$$(x-2)(y-1) = 3$$

$$y-1 = \frac{3}{x-2}$$


$$y = \frac{3}{x-2} + 1$$

So  $f^{-1}(x) = \frac{3}{x-2} + 1$

**(1 mark)**

b.  $d_{f^{-1}} = r_f$

Do a quick sketch of  $y = \frac{3}{x-1} + 2$



$$r_f = (2, \infty)$$

So  $d_{f^{-1}} = (2, \infty)$

**(1 mark)**

**Question 4**

a.  $f(x) = \sin(e^{2x})$

Let  $y = \sin(e^{2x})$

$$y = \sin(u)$$

$$\frac{dy}{du} = \cos(u)$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} \text{ (Chain rule)}$$

$$= \cos(u) \cdot 2e^{2x}$$

$$= 2e^{2x} \cos(e^{2x})$$

$$\text{Let } u = e^{2x}$$

$$\frac{du}{dx} = 2e^{2x}$$

**(1 mark)** – for knowing to use the chain rule  
and attempting to use it

**(1 mark)** – correct answer

b.

$$y = \frac{\log_e(x)}{x^3 + 2x}$$

$$\frac{dy}{dx} = \frac{(x^3 + 2x) \times \frac{1}{x} - (3x^2 + 2) \log_e(x)}{(x^3 + 2x)^2}$$

$$= \frac{x^2 + 2 - (3x^2 + 2) \log_e(x)}{(x^3 + 2x)^2}$$

**(1 mark)** – for knowing to use the quotient rule  
and attempting to use it

**(1 mark)** – correct answer

**Question 5**

For  $y = 2 \sin\left(2\left(x - \frac{\pi}{3}\right)\right)$  the period is  $\frac{2\pi}{2} = \pi$  **(1 mark)** correct period

This means that for  $x \in [-\pi, \pi]$  there will be two complete periods of the graph.

x-interceptsMethod 1

The graph is a sine graph with a period of  $\pi$  that has been translated  $\frac{\pi}{3}$  units to the right so

that the x-intercepts will occur at  $\left(\frac{\pi}{3}, 0\right)$ , at  $\left(\frac{\pi}{2} + \frac{\pi}{3}, 0\right) = \left(\frac{5\pi}{6}, 0\right)$ , at

$\left(-\frac{\pi}{2} + \frac{\pi}{3}, 0\right) = \left(-\frac{\pi}{6}, 0\right)$  and at  $\left(-\pi + \frac{\pi}{3}, 0\right) = \left(-\frac{2\pi}{3}, 0\right)$ . **(1 mark)** for x- intercepts

Method 2

The x- intercepts occurs when  $y = 0$ .

$$y = 2 \sin\left(2\left(x - \frac{\pi}{3}\right)\right)$$

becomes

$$0 = 2 \sin\left(2\left(x - \frac{\pi}{3}\right)\right)$$

So,  $\sin\left(2\left(x - \frac{\pi}{3}\right)\right) = 0$

$$2\left(x - \frac{\pi}{3}\right) = \dots - 3\pi, -2\pi, -\pi, 0, \pi, 2\pi, \dots$$

$$x - \frac{\pi}{3} = \dots -\frac{3\pi}{2}, -\pi, -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi, \dots$$

$$x = \dots -\frac{7\pi}{6}, -\frac{2\pi}{3}, -\frac{\pi}{6}, \frac{\pi}{3}, \frac{5\pi}{6}, \frac{4\pi}{3}, \dots$$

$$x = \frac{-2\pi}{3}, -\frac{\pi}{6}, \frac{\pi}{3}, \frac{5\pi}{6} \text{ since } -\pi \leq x \leq \pi$$

**(1 mark)** for x- intercepts

y - intercept ( $x = 0$ )

$$y = 2 \sin\left(2\left(0 - \frac{\pi}{3}\right)\right)$$

$$= 2 \sin\left(-\frac{2\pi}{3}\right)$$


$$= 2 \times -\sin\left(\frac{\pi}{3}\right)$$

$$= 2 \times -\frac{\sqrt{3}}{2}$$

$$= -\sqrt{3}$$

y-intercept occurs at  $(0, -\sqrt{3})$

**(1 mark)**



Angle measured clockwise because it is negative.

To find the endpoints:

Method 1

Because two complete periods occur for  $x \in [-\pi, \pi]$  the right endpoint and left endpoint will have the same  $y$ -coordinate and it will be the same as the  $y$ -intercept.

So left endpoint is  $(-\pi, -\sqrt{3})$  and the right endpoint is  $(\pi, -\sqrt{3})$ .

**(1 mark)**

Method 2


$$\begin{aligned} \text{Left endpoint: } f(-\pi) &= 2 \sin\left(2\left(-\pi - \frac{\pi}{3}\right)\right) & \text{Right endpoint: } f(\pi) &= 2 \sin\left(2\left(\pi - \frac{\pi}{3}\right)\right) \\ &= 2 \sin\left(2\left(\frac{-4\pi}{3}\right)\right) & &= 2 \sin\left(2\left(\frac{2\pi}{3}\right)\right) \\ &= 2 \sin\left(\frac{-8\pi}{3}\right) & &= 2 \sin\left(\frac{4\pi}{3}\right) \\ &= 2 \sin\left(\frac{-2\pi}{3}\right) & &= 2 \times -\sin\left(\frac{\pi}{3}\right) \\ &= 2 \times -\sin\left(\frac{\pi}{3}\right) & &= 2 \times -\frac{\sqrt{3}}{2} \\ &= 2 \times -\frac{\sqrt{3}}{2} & &= -\sqrt{3} \\ &= -\sqrt{3} \end{aligned}$$

Left endpoint is  $(-\pi, -\sqrt{3})$ .

Right endpoint is  $(\pi, -\sqrt{3})$ .

**(1 mark)**


The graph of  $y = 2 \sin\left(2\left(x - \frac{\pi}{3}\right)\right)$  is shown below.



**(1 mark)** correct shape including amplitude

## Question 6

a.



$$\begin{aligned}
 \Pr(X > 9) &= \Pr(X < 5) \text{ by symmetry} \\
 &= \Pr(Z < -1) \\
 &= 0.16 \text{ (given)}
 \end{aligned}$$

(1 mark)

b.  $\Pr(X < 7 | X < 9)$  (conditional probability)

$$= \frac{\Pr(X < 7 \cap X < 9)}{\Pr(X < 9)}$$

(1 mark)

$$= \frac{\Pr(X < 7)}{1 - \Pr(X > 9)}$$

$$= \frac{0.5}{1 - 0.16} \text{ from part a.}$$

(1 mark)

$$= \frac{0.5}{0.84}$$

$$= \frac{50}{84}$$

$$= \frac{25}{42}$$

(1 mark)

**Question 7**

- a. Since  $f(x)$  represents a probability density function then

$$\int_1^3 (ax+1)dx = 1 \quad (1 \text{ mark})$$

$$\left[ \frac{ax^2}{2} + x \right]_1^3 = 1$$

$$\left\{ \left( \frac{9a}{2} + 3 \right) - \left( \frac{a}{2} + 1 \right) \right\} = 1$$

$$\frac{8a}{2} + 2 = 1$$

$$4a = -1$$

$$a = -\frac{1}{4} \text{ as required}$$

**(1 mark)**

- b.

$$\Pr(X < 2) = \int_1^2 \left( -\frac{1}{4}x + 1 \right) dx \quad (1 \text{ mark})$$

$$= \left[ -\frac{1}{4} \times \frac{x^2}{2} + x \right]_1^2$$


$$= \left[ -\frac{x^2}{8} + x \right]_1^2$$

$$= \left\{ \left( -\frac{4}{8} + 2 \right) - \left( -\frac{1}{8} + 1 \right) \right\}$$

$$= -\frac{3}{8} + 1$$

$$= \frac{5}{8}$$

**(1 mark)**

**Question 8****a.**

Note that at the points  $(1,0)$ ,  $(3,0)$  and  $(5,0)$  the graph of  $y = |f(x)|$  has cusps; i.e. “pointy bits” not smooth curves since the graph of  $y = f(x)$  is being reflected in the  $x$ -axis.

**(1 mark)**

**b.** 
$$A = \int_1^3 f(x)dx - \int_3^5 f(x)dx$$

**(1 mark)**



**Question 9****a.**

$$\begin{aligned}
 \text{area} &= \frac{1}{2} \times \text{base} \times \text{height} \\
 &= \frac{1}{2} \times a \times b \\
 &= \frac{1}{2} \times a \times f(a) \text{ where } f(x) = 1 - \frac{x}{5} \\
 &= \frac{1}{2} \times a \times \left(1 - \frac{a}{5}\right) \\
 &= \frac{a}{2} - \frac{a^2}{10} \text{ as required}
 \end{aligned}$$

**(1 mark)****b.**

$$\begin{aligned}
 \text{Let } A &= \frac{a}{2} - \frac{a^2}{10} \\
 \frac{dA}{da} &= \frac{1}{2} - \frac{2a}{10}
 \end{aligned}$$

**(1 mark)**

Maximum occurs when  $\frac{dA}{da} = 0$

$$\frac{1}{2} - \frac{a}{5} = 0$$

$$\frac{a}{5} = \frac{1}{2}$$

$$2a = 5$$

$$a = \frac{5}{2}$$

Note: we know we have a maximum because the graph of the function  $A = \frac{a}{2} - \frac{a^2}{10}$  is an inverted parabola.

We have a maximum when  $a = \frac{5}{2}$ .

**(1 mark)**

$$\begin{aligned}
 \text{When } a = \frac{5}{2}, \text{ area} &= \frac{a}{2} - \frac{a^2}{10} \\
 &= \frac{5}{4} - \frac{25}{40} \\
 &= \frac{50}{40} - \frac{25}{40} \\
 &= \frac{25}{40} \\
 &= \frac{5}{8} \text{ square units}
 \end{aligned}$$

$$\begin{aligned}
 \text{or when } a = 2.5, \text{ area} &= \frac{2.5}{2} - \frac{2.5^2}{10} \\
 &= 1.25 - \frac{6.25}{10} \\
 &= 1.25 - 0.625 \\
 &= 0.625 \text{ square units}
 \end{aligned}$$

**(1 mark)**

**Question 10**

$$y = 3x^2 + a$$

$$\frac{dy}{dx} = 6x$$

The gradient of a normal to  $y = 3x^2 + a$  is  $-\frac{1}{6x}$ .

**(1 mark)**

Also the gradient of the normal  $y = \frac{x}{3} + 1$  is  $\frac{1}{3}$ .

$$\begin{aligned} \text{When } -\frac{1}{6x} &= \frac{1}{3} \\ -3 &= 6x \\ x &= -\frac{1}{2} \end{aligned}$$

**(1 mark)**

The  $x$ -coordinate of the point where the normal hits the curve is  $-\frac{1}{2}$ .

$$\begin{aligned} \text{So } y &= -\frac{1}{2} \div 3 + 1 \\ &= -\frac{1}{2} \times \frac{1}{3} + 1 \\ &= -\frac{1}{6} + 1 \\ &= \frac{5}{6} \end{aligned}$$

The curve and the normal both pass through the point  $\left(-\frac{1}{2}, \frac{5}{6}\right)$ .

Substituting this point into

**(1 mark)**

$$\begin{aligned} y &= 3x^2 + a \\ \text{gives } \frac{5}{6} &= 3 \times \left(-\frac{1}{2}\right)^2 + a \\ \frac{5}{6} &= 3 \times \frac{1}{4} + a \\ \text{So } a &= \frac{5}{6} - \frac{3}{4} \\ &= \frac{10 - 9}{12} \\ &= \frac{1}{12} \end{aligned}$$

**(1 mark)**

**Question 11**

a.  $y = e^{2x}$

$$y = 6 - e^x$$

At the point of intersection of the graphs,

$$e^{2x} = 6 - e^x$$

$$e^{2x} + e^x - 6 = 0$$

Let  $e^x = m$

$$m^2 + m - 6 = 0$$

**(1 mark)**

$$(m + 3)(m - 2) = 0$$

$$m = -3 \text{ or } m = 2$$

So  $e^x = -3$  or  $e^x = 2$

no solution  $x = \log_e(2)$

The  $x$ -coordinate of the point of intersection is  $\log_e(2)$ .

**(1 mark)**

b.

$$\text{Area} = \int_0^{\log_e(2)} \{(6 - e^x) - (e^{2x})\} dx$$

**(1 mark)**

$$= \int_0^{\log_e(2)} (6 - e^x - e^{2x}) dx$$

$$= \left[ 6x - e^x - \frac{e^{2x}}{2} \right]_0^{\log_e(2)}$$

**(1 mark)**

$$= \left\{ \left( 6 \log_e(2) - e^{\log_e(2)} - \frac{e^{2 \log_e(2)}}{2} \right) - \left( 0 - e^0 - \frac{e^0}{2} \right) \right\}$$

$$= \log_e(2^6) - 2 - \frac{e^{\log_e(2^2)}}{2} + 1 + \frac{1}{2}$$

**(1 mark)**

$$= \log_e(2^6) - 2 - \frac{4}{2} + 1 + \frac{1}{2}$$

$$= \log_e(64) - \frac{5}{2} \text{ square units}$$

**(1 mark)****Total 40 marks**