

# MATHEMATICAL METHODS (CAS)

## Units 3 & 4 – Written examination 1



### 2012 Trial Examination

### **SOLUTIONS**

#### Question 1

a.  $f(x^4 + 2) = \log_e(x^4 + 2 + 3) = \log_e(x^4 + 5)$

M1+A1  
2 marks

b.  $f'(g(x)) = \frac{1}{(x^4+5)} \times 4x^3 = \frac{4x^3}{(x^4+5)}$

M1+A1  
2 marks

c.  $f'(g(-2)) = \frac{4(-2)^3}{((-2)^4+5)} = \frac{-32}{21}$

A1  
1 mark

#### Question 2

a.  $\left(\frac{5x^2}{2} - 10x\right)_0^a = 0$

$$\begin{aligned}\frac{5a^2}{2} - 10a &= 0 \\ 5a(a - 4) &= 0 \\ a = 0 \text{ or } a &= 4\end{aligned}$$

Since  $a \neq 0$ ,  $a = 4$

M1+A1  
2 marks

2012 MATHMETH(CAS) EXAM 1

b.  $\int_0^\pi \cos\left(\frac{x}{2}\right) dx = 2\sin\left(\frac{x}{2}\right)_0^\pi = 2$

M1+A1  
2 marks

**Question 3**

a.  $x = -4e^{\frac{y}{2}} + 1$

$$\begin{aligned} \frac{x-1}{-4} &= e^{\frac{y}{2}} \\ \frac{y}{2} &= \log_e\left(\frac{1-x}{4}\right) \\ f^{-1}(x) &= 2\log_e\left(\frac{1-x}{4}\right) \end{aligned}$$

M2+A1  
3 marks

b.  $1-x > 0$

Domain of  $f^{-1}(x)$  is  $(-\infty, 1)$

M1  
1 mark

c.  $2\log_e\left(\frac{1-x}{4}\right) = 0$

$$\begin{aligned} \left(\frac{1-x}{4}\right) &= 1 \\ \frac{1-x}{4} &= 1 \\ x &= -3 \end{aligned}$$

M1+A1  
2 marks

**Question 4**

a.  $3\cos(2x) = -\frac{3\sqrt{3}}{2}$

$$\begin{aligned} \cos(2x) &= -\frac{\sqrt{3}}{2} \quad \text{for } -\pi \leq 2x \leq \pi \\ 2x &= \frac{5\pi}{6}, -\frac{5\pi}{6} \\ x &= \frac{5\pi}{12}, -\frac{5\pi}{12} \end{aligned}$$

M1+A1  
2 marks

2012 MATHMETH(CAS) EXAM 1

b.  $-3 \sin(2x) \times 2 = 0$

$$2x = 0, \pi, -\pi$$

$$x = 0, \frac{\pi}{2}, -\frac{\pi}{2}$$

M1+A1

2 marks

c.  $(0, \frac{\pi}{2})$

M1

1 mark

**Question 5**

$$\begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix} \left( \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -1 \\ 5 \end{bmatrix} \right) = \begin{bmatrix} x' \\ y' \end{bmatrix}$$

$$\begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x - 1 \\ y + 5 \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix}$$

$$\begin{bmatrix} -x + 1 \\ -2y - 10 \end{bmatrix} = \begin{bmatrix} x' \\ y' \end{bmatrix}$$

$$-x + 1 = x' \text{ and } -2y - 10 = y'$$

$$x = -x' \mp 1 \text{ and } y = \frac{-y' - 10}{2}$$

$$\frac{-y' - 10}{2} = |-x' \mp 1 + 2| - 3$$

$$y' + 10 = -2|x' - 3| + 6$$

$$y = -2|x - 3| - 4$$

M2+A2

4 marks

**Question 6**

a.  $V = \frac{1}{3}\pi r^2 h = 18\pi$

M1

1 mark

b.  $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi \left(\frac{h}{2}\right)^2 \times h = \frac{\pi}{12}h^3$

$$\frac{dh}{dt} = \frac{dh}{dV} \times \frac{dV}{dt} = \frac{4}{\pi h^2} \times 2 = \frac{2}{9\pi} m/s$$

M1+A1

2 marks

**Question 7**

$$\begin{aligned}f(x) &= \sqrt{x}, \quad f'(x) = \frac{1}{2\sqrt{x}} \\ \sqrt{24.95} &\approx \sqrt{25} - 0.05 \times \frac{1}{2\sqrt{25}} \\ &\approx 5 - \frac{0.05}{10} \approx 4.995\end{aligned}$$

M1+A2  
3 marks

**Question 8**

a.  $\int_1^4 k(-x^2 + 5x - 4)dx = 1$

$$k \left( -\frac{x^3}{3} + \frac{5x^2}{2} - 4x \right)_1^4 = 1$$

which gives  $k = \frac{2}{9}$

M1+A2  
3 marks

b.  $\Pr(X > 3) = \frac{2}{9} \int_3^4 (-x^2 + 5x - 4)dx$

$$\begin{aligned}&= \frac{2}{9} \left( \frac{-64}{3} + 40 - 16 + \frac{27}{3} - \frac{45}{2} + 12 \right) \\&= \frac{7}{27}\end{aligned}$$

A2  
2 marks

**Question 9**

a.  $m = 200$  (95% means 2 standard deviations from the mean)

M1  
1 mark

b.  $\Pr(X < 210) = \Pr \left( Z < \frac{210-220}{10} \right)$

$$= \Pr(Z < -1) = \Pr(Z > 1) = 0.16$$

M1+A1  
2 marks

c.  $\Pr(X > 230 | X > 220) = \frac{\Pr(X > 230)}{\Pr(X > 220)} = \frac{\Pr(Z > 1)}{\Pr(Z > 0)} = \frac{0.16}{0.5} = \frac{16}{50} = \frac{8}{25} \text{ or } 0.32$

M1+A1  
2 marks