

MAV Specialist Mathematics Examination 1

Answers & Solutions

Part I (Multiple-choice) Answers

- | | | | | |
|-------|-------|-------|-------|-------|
| 1. E | 2. B | 3. C | 4. A | 5. C |
| 6. A | 7. A | 8. B | 9. D | 10. D |
| 11. A | 12. B | 13. C | 14. C | 15. D |
| 16. A | 17. E | 18. A | 19. E | 20. E |
| 21. E | 22. D | 23. C | 24. B | 25. E |
| 26. E | 27. D | 28. E | 29. B | 30. D |

Question 1

[E]

$$\begin{aligned}\vec{AB} &= \vec{AO} + \vec{OB} \\ &= \vec{OB} - \vec{OA} \\ &= (4\vec{i} - 2\vec{j}) - (5\vec{i} - 2\vec{j} - \vec{k}) \\ &= -\vec{i} + \vec{k}\end{aligned}$$

Question 2

[B]

$$\begin{aligned}|z - 1 + 2i| &= 2 \\ |x + yi - 1 + 2i| &= 2 \\ |(x - 1) + (y + 2)i| &= 2 \\ \sqrt{(x - 1)^2 + (y + 2)^2} &= 2 \\ (x - 1)^2 + (y + 2)^2 &= 4 \\ \text{circle, centre } (1, -2), \text{ radius } 2\end{aligned}$$

Question 3

[C]

$$\begin{aligned}-1 \leq 3x - 1 \leq 1 & \quad 3x - 1 \leq 1 \\ 3x - 1 \geq -1 & \quad 3x \leq 2 \\ 3x \geq 0 & \quad x \leq \frac{2}{3} \\ x \geq 0\end{aligned}$$

OR

$$-2\sin^{-1}(3x - 1) - 2 = -2\sin^{-1}\left[3\left(x - \frac{1}{3}\right)\right] - 2$$

$$f(x) = \sin^{-1}(x), \text{ dom } f = [-1, 1]$$

$$g(x) = \sin^{-1}(3x) \quad \text{Dilated by a factor of } \frac{1}{3}.$$

$$\text{dom } g = \left[-\frac{1}{3}, \frac{1}{3}\right]$$

$$h(x) = \sin^{-1}\left[3\left(x - \frac{1}{3}\right)\right] \text{ is } g(x) \text{ translated } \frac{1}{3}$$

units right.

$$\text{dom } h = \left[0, \frac{2}{3}\right]$$

Note: The two (-2)'s in the given function do not effect the domain.

Question 4


[A]

$$\begin{aligned}\sec^2 x &= 1 + \tan^2 x \\ &= 1 + \frac{4}{25} \\ &= \frac{29}{25}\end{aligned}$$

$$\sec x = -\frac{\sqrt{29}}{5}, \text{ since } \pi < x < \frac{3\pi}{2}$$

Question 5

[C]



$$\tan \theta = \frac{-1}{-1} = 1$$

$$\theta = \frac{5\pi}{4}$$

$$\therefore -1 - i = \sqrt{2} \operatorname{cis}\left(-\frac{3\pi}{4}\right)$$

Question 6

[A]

$$\begin{aligned} \int \frac{3}{\sqrt{1-4x^2}} dx &= -3 \int \frac{-1}{\sqrt{4\left(\frac{1}{4} - x^2\right)}} dx \\ &= -3 \int \frac{-1}{2\sqrt{\left(\frac{1}{2}\right)^2 - x^2}} dx \\ &= -\frac{3}{2} \operatorname{Cos}^{-1}(2x) + c \end{aligned}$$

Question 7

[A]

$$\begin{aligned} \int_2^3 \frac{4}{x^2} \log_e\left(\frac{2}{x}\right) dx & \quad \text{Let } u = \frac{2}{x} = 2x^{-1} \\ \frac{du}{dx} &= -2x^{-2} = \frac{-2}{x^2} \\ -2 \frac{du}{dx} &= \frac{4}{x^2} \end{aligned}$$

$$\begin{aligned} \text{Terminals: } x = 3, u &= \frac{2}{3} \\ x = 2, u &= 1 \end{aligned}$$

$$-2 \int_1^{\frac{2}{3}} \log_e(u) du$$

Question 8

[B]

$$\begin{aligned} (1+i)^4 &= \left(\sqrt{2} \operatorname{cis}\left(\frac{\pi}{4}\right)\right)^4 \\ &= 4 \operatorname{cis}(\pi) \\ &= 4(\cos(\pi) + i \sin(\pi)) \\ &= -4 \end{aligned}$$

Question 9

[D]

$$\text{If } z = r \operatorname{cis}(\theta), \text{ then } \bar{z} = r \operatorname{cis}(-\theta)$$

Question 10

[D]

$$\begin{aligned} \tilde{r}(t) &= 3e^{-t} \tilde{i} + \frac{3}{2} \cos(2t) \tilde{j} \\ \dot{\tilde{r}}(t) &= -3e^{-t} \tilde{i} - \frac{3}{2} \times 3 \sin(2t) \tilde{j} \\ \dot{\tilde{r}}(0) &= -3\tilde{i} \\ \left| \dot{\tilde{r}}(0) \right| &= 3 \\ &= 3 \end{aligned}$$

Question 11

[A]

$$\begin{aligned} \text{Area} &= \frac{1}{2} \left(1 + \frac{8}{9}\right) \frac{1}{2} + \frac{1}{2} \left(\frac{8}{9} + \frac{1}{2}\right) \frac{1}{2} \\ &= \frac{1}{4} \left(\frac{17}{9} + \frac{25}{18}\right) \\ &= \frac{1}{4} \left(\frac{59}{18}\right) \\ &= \frac{59}{72} \end{aligned}$$

Question 12**[B]**

$$\begin{aligned} & \int \frac{1}{x \log_e(5x)} dx \\ &= \int \frac{\frac{1}{x}}{\log_e(5x)} dx \quad \left[= \int \frac{f'(x)}{f(x)} dx \right] \\ &= \log_e(\log_e(5x)) + c \end{aligned}$$

OR

In order to determine the correct response to this question, students could systematically differentiate the options.

$$\text{If } y = \log_e(5x), \quad \frac{dy}{dx} = \frac{1}{x}$$

$$\text{If } y = \log_e(\log_e(5x)), \quad \text{Let } u = \log_e(5x)$$

$$y = \log_e(u) \quad \frac{du}{dx} = \frac{1}{x}$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \times \frac{du}{dx} \\ &= \frac{1}{u} \times \frac{1}{x} \\ &= \frac{1}{x \log_e(5x)} \end{aligned}$$

Question 13**[C]**

Before translation, asymptotes given by

$$y = \pm \frac{b}{a}x \quad \text{or use } (y - k) = \pm \frac{b}{a}(x - h)$$

$$y = \pm \frac{6}{3}(x - 2)$$

$$y = \pm 2(x - 2)$$

Question 14**[C]**

$$\begin{aligned} \int_1^4 \frac{x-2}{x^2-4x+7} dx &= \frac{1}{2} \int_1^4 \frac{2x-4}{x^2-4x+7} dx \\ &= \frac{1}{2} \left[\log_e(x^2-4x+7) \right]_1^4 \\ &= \frac{1}{2} (\log_e 7 - \log_e 4) \\ &= \frac{1}{2} \log_e \left(\frac{7}{4} \right) \\ &= \log_e \left(\sqrt{\frac{7}{4}} \right) \\ &= \log_e \left(\frac{\sqrt{7}}{2} \right) \end{aligned}$$

Question 15**[D]**

$$a = 2i - 3j + 4k, \quad |a| = \sqrt{16 + 9 + 4} = \sqrt{29}$$

$$b = 3i + 2j + k, \quad |b| = \sqrt{9 + 4 + 1} = \sqrt{14}$$

$$\begin{aligned} \cos \theta &= \frac{a \cdot b}{|a| \times |b|} \\ &= \frac{2(3) - 3(2) + 4(1)}{\sqrt{29} \times \sqrt{14}} \\ &= \frac{4}{\sqrt{406}} \end{aligned}$$

$$\theta = \text{Cos}^{-1} \left(\frac{4}{\sqrt{406}} \right) = 78.50^\circ$$

Question 16**[A]**

If l grams dissolved, then $(8 - l)$ grams undissolved.

$$\begin{aligned} \text{Hence } \frac{dl}{dt} &= \frac{25}{100}(8 - l) \\ &= \frac{8 - l}{4} \end{aligned}$$

Question 17**[E]**

$$y = \sin(3x - 1)$$

$$\frac{dy}{dx} = 3 \cos(3x - 1)$$

$$\frac{d^2y}{dx^2} = -9 \sin(3x - 1)$$

$$\begin{aligned} 9y + 3 \frac{dy}{dx} + \frac{d^2y}{dx^2} \\ &= 9 \sin(3x - 1) + 9 \cos(3x - 1) - 9 \sin(3x - 1) \\ &= 9 \cos(3x - 1) \end{aligned}$$

Question 18

[A]

$$\begin{aligned}
 uv &= 3 \times 2cis\left(\frac{5\pi}{6} + \frac{3\pi}{4}\right) \\
 &= 6cis\left(\frac{10\pi + 9\pi}{12}\right) \\
 &= 6cis\left(\frac{19\pi}{12}\right) \\
 &= 6cis\left(-\frac{5\pi}{12}\right) \text{ since } -\pi < \theta < \pi
 \end{aligned}$$

Question 19

[E]

Method 1: Consider total motion.

$$s = 40\text{m}, a = 9.8\text{m/s}^2, u = -15\text{m/s}$$

$$s = ut + \frac{1}{2}at^2$$

$$40 = -15t + 4.9t^2$$

$$4.9t^2 - 15t - 40 = 0$$

 Quadratic formula $\Rightarrow t = 4.77\text{sec}$, since $t > 0$

Method 2: Consider upwards and downwards motions separately.

Upward motion:

$$u = 15\text{m/s}, v = 0\text{m/s}, a = -9.8\text{m/s}^2$$

$$t = \frac{v - u}{a} = \frac{-15}{-9.8} = 1.53 \text{ seconds}$$

$$v^2 - u^2 = 2as$$

$$s = \frac{-u^2}{2a} = \frac{-15^2}{-19.6} = 11.48$$

Downward motion:

$$s = 40 + 11.48 = 51.48 \text{ metres,}$$

$$u = 0\text{m/s}, a = 9.8\text{m/s}^2$$

$$s = ut + \frac{1}{2}at^2$$

$$51.48 = 4.9t^2$$

$$t = 3.24 \text{ seconds, since } t > 0$$

 Total time = $1.53 + 3.24 = 4.77$ seconds

Question 20

[E]

$$\frac{dy}{dx} = \frac{1}{x^2}, \quad y_{n+1} = y_n + hf'(x_n), \quad h = 0.2$$

x	y
1	3
1.2	$3 + 0.2\left(\frac{1}{1^2}\right) = 3.2$
1.4	$3.2 + 0.2\left(\frac{1}{1.2^2}\right) = 3.339$

Question 21

[E]

$$\begin{aligned}
 \left| -2\underset{\sim}{i} + 6\underset{\sim}{j} - 9\underset{\sim}{k} \right| &= \sqrt{4 + 36 + 81} \\
 &= 11
 \end{aligned}$$

$$\begin{aligned}
 \frac{22}{11} \left(-2\underset{\sim}{i} + 6\underset{\sim}{j} - 9\underset{\sim}{k} \right) \\
 &= 2 \left(-2\underset{\sim}{i} + 6\underset{\sim}{j} - 9\underset{\sim}{k} \right) \\
 &= -4\underset{\sim}{i} + 12\underset{\sim}{j} - 18\underset{\sim}{k}
 \end{aligned}$$

$$\text{Option E: } 4\underset{\sim}{i} - 12\underset{\sim}{j} + 18\underset{\sim}{k} = -2(-2\underset{\sim}{i} + 6\underset{\sim}{j} - 9\underset{\sim}{k})$$

Question 22

[D]

$$\begin{aligned}
 \left| \underset{\sim}{b} \right| &= \sqrt{5} \quad \underset{\sim}{a} \cdot \underset{\sim}{b} = \frac{1}{\sqrt{5}} ((2 \times 1) + (-3 \times -2)) \\
 &= \frac{8}{\sqrt{5}}
 \end{aligned}$$

Question 23

[C]

$$\begin{aligned}
 a &= v \frac{dv}{dx}, \quad \frac{dv}{dx} = \frac{1}{2} \\
 &= \frac{x}{2} \times \frac{1}{2} \\
 &= \frac{x}{4}
 \end{aligned}$$

Question 24

$$\begin{aligned} \vec{r} &= \vec{i} - 2\vec{j} + \vec{k} \\ \vec{r} &= t\vec{i} - 2t\vec{j} + t\vec{k} + \vec{c} \\ t = 0, \vec{r} &= 2\vec{i} - 3\vec{k}, \Rightarrow \vec{c} = 2\vec{i} - 3\vec{k} \\ \vec{r} &= (t+2)\vec{i} - 2t\vec{j} + (t-3)\vec{k} \end{aligned}$$

Question 25


$$\begin{aligned} y &= \sin^{-1}\left(\frac{2}{x}\right) \quad \text{Let } u = \frac{2}{x} = 2x^{-1} \\ \frac{du}{dx} &= -2x^{-2} = -\frac{2}{x^2} \\ \frac{dy}{dx} &= \frac{dy}{du} \times \frac{du}{dx} \\ &= \frac{1}{\sqrt{1-u^2}} \times \frac{-2}{x^2} \\ &= \frac{-2}{x^2 \sqrt{1-\frac{4}{x^2}}} \\ &= \frac{-2}{x^2 \sqrt{\frac{x^2-4}{x^2}}} \\ &= \frac{-2}{x^2 \frac{\sqrt{x^2-4}}{x}} \\ &= \frac{-2}{x\sqrt{x^2-4}} \end{aligned}$$

Question 26

$$\begin{aligned} \frac{dv}{dt} &= 3e^{-0.2t} + 2 \\ v &= \int (3e^{-0.2t} + 2) dt \\ v &= -15e^{-0.2t} + 2t + c \\ \text{Starts from rest } (t = 0, v = 0), c &= 15 \\ \text{Hence } v &= -15e^{-0.2t} + 2t + 15 \\ \text{When } t = 2, v &= 8.95 \end{aligned}$$

[B]

Question 27



[D]

[E]

$$\begin{aligned} N &= mg \cos \theta, \quad F = \mu N = \mu mg \cos \theta \\ R &= mg \sin \theta - F, \\ &= mg \sin \theta - \mu mg \cos \theta \\ a &= \frac{R}{m} = g \sin \theta - \mu g \cos \theta \end{aligned}$$

Question 28

$$\begin{aligned} R &= N - 4g \\ N - 4g &= 4 \times 3 \\ N &= 12 + 4g \\ &= 51.2 \end{aligned}$$

[E]

Question 29

$$\begin{aligned} \Delta p &= p_{\text{final}} - p_{\text{initial}}, \quad p = m \times v \\ &= 4(5) - 4(8) \\ &= -12 \end{aligned}$$

[B]

[E]

Question 30

$$\begin{aligned} \text{Long diagonal: } \vec{AC} &= \vec{AO} + \vec{OD} \\ &= \vec{OD} - \vec{OA} \\ &= \vec{d} - \vec{a} \end{aligned}$$

$$\text{Short diagonal: } \vec{OB} = \vec{b}$$

$$\text{Diagonals are perpendicular if } (\vec{d} - \vec{a}) \cdot \vec{b} = 0$$

[D]

Part II Solutions

Question 1

$$\begin{aligned} \text{a. } \frac{1}{9-x^2} &\equiv \frac{a}{3-x} + \frac{b}{3+x} \\ &\equiv \frac{a(3+x) + b(3-x)}{(3-x)(3+x)} \\ \therefore 1 &\equiv a(3+x) + b(3-x) \quad [\text{A}] \end{aligned}$$

$$\text{If } x = 3, 1 = 6a, a = \frac{1}{6}$$

$$\text{If } x = -3, 1 = 6b, b = \frac{1}{6}$$

$$\therefore \frac{1}{9-x^2} = \frac{1}{6(3-x)} + \frac{1}{6(3+x)} \quad [\text{A}]$$

$$\text{b. } \int \frac{2}{\sqrt{9-x^2}} dx = 2\text{Sin}^{-1}\left(\frac{x}{3}\right) \quad [\text{A}]$$

$$\text{c. } V = \int_a^b (\pi y^2) dx$$

$$y = \frac{1}{\sqrt{9-x^2}} + 1$$

$$y^2 = \frac{1}{9-x^2} + \frac{2}{\sqrt{9-x^2}} + 1$$

$$V = \pi \int_0^2 \left(\frac{1}{9-x^2} + \frac{2}{\sqrt{9-x^2}} + 1 \right) dx \quad [\text{M}]$$

$$= \pi \int_0^2 \left\{ \frac{1}{6} \left(\frac{1}{3-x} + \frac{1}{3+x} \right) + \frac{2}{\sqrt{9-x^2}} + 1 \right\} dx$$

$$= \pi \left[\frac{1}{6} \log_e \left(\frac{3+x}{3-x} \right) + 2\text{Sin}^{-1}\left(\frac{x}{3}\right) + x \right]_0^2 \quad [\text{M}]$$

$$= \pi \left[\left(\frac{1}{6} \log_e 5 + 2\text{Sin}^{-1} \frac{2}{3} + 2 \right) - 0 \right]$$

$$\approx 11.71 \text{ cubic units} \quad [\text{A}]$$

Question 2

$$h'(x) = \frac{x}{\sqrt{1-x}} \quad \text{Let } u = 1-x$$

$$\frac{du}{dx} = -1$$

$$x = 1-u$$

$$h(x) = -\int \frac{1-u}{\frac{1}{u^2}} du = \int (-u^{-\frac{1}{2}} + u^{\frac{1}{2}}) du \quad [\text{M}]$$

$$= -2u^{\frac{1}{2}} + \frac{2}{3}u^{\frac{3}{2}} + c$$

$$= \frac{2}{3}(1-x)\sqrt{1-x} - 2\sqrt{1-x} + c \quad [\text{M}]$$

$$= \frac{2}{3}\sqrt{1-x}(1-x-3) + c$$

$$= -\frac{2}{3}\sqrt{1-x}(x+2) + c$$

$$\text{Since } h(-3) = \frac{4}{3},$$

$$\frac{4}{3} = -\frac{2}{3}(-1)\sqrt{4} + c, c = 0 \quad [\text{A}]$$

$$h(x) = -\frac{2}{3}(x+2)\sqrt{1-x}$$

Question 3

$$\text{a. } w = 1 - \sqrt{3}i$$

$$\text{Arg } w = -\frac{\pi}{3} \quad [\text{A}]$$

$$\text{b. } \arg(z^2w) = \left(2 \times \frac{3\pi}{4} \right) - \frac{\pi}{3}$$

$$= \frac{3\pi}{2} - \frac{\pi}{3}$$

$$= \frac{7\pi}{6}$$


$$\text{Arg}(z^2w) = -\frac{5\pi}{6} \quad [\text{A}]$$

$$\text{c. } \frac{z}{w} = \frac{3}{2} \text{cis} \left(\frac{3\pi}{4} - \left(-\frac{\pi}{3} \right) \right)$$

$$= \frac{3}{2} \text{cis} \left(\frac{9\pi}{12} + \frac{4\pi}{12} \right)$$

$$= \frac{3}{2} \text{cis} \left(\frac{13\pi}{12} \right)$$

$$= \frac{3}{2} \text{cis} \left(-\frac{11\pi}{12} \right) \quad [\text{A}]$$

Question 4


- a. $R^2 = 3^2 + 4^2 - 2(4)(3)\cos(105^\circ)$
 Diagram [M]
 $R \approx 5.59$ Newtons [A]

- b. Let the angle between the 3 Newton force and the Resultant be θ .


$$\text{By sine rule } \frac{5.59}{\sin(105^\circ)} = \frac{4}{\sin \theta} \quad [\text{M}]$$

$$\sin \theta = \frac{4 \sin(105^\circ)}{5.59}$$

$$\theta = 43.755^\circ$$

Hence angle between the Resultant and North is $43.755^\circ - 30^\circ = 13.755^\circ$

$$a = \frac{F}{m} = 2.79 \text{ m/s}^2$$



$$\sin(13.755^\circ) = \frac{c}{2.79} \Leftrightarrow c = 2.79 \sin(13.755^\circ)$$

[M]

$$c = 0.664$$

$$\cos(13.755^\circ) = \frac{d}{2.79} \Leftrightarrow d = 2.79 \cos(13.755^\circ)$$

$$d = 2.71$$

$$\text{Hence } \underline{\underline{a}} = -0.66 \underline{\underline{i}} + 2.71 \underline{\underline{j}} \quad [\text{A}]$$

OR



$$\underline{\underline{\Sigma F}} = 3 \cos 60^\circ \underline{\underline{i}} + 3 \sin 60^\circ \underline{\underline{j}} - 4 \cos 45^\circ \underline{\underline{i}} + 4 \sin 45^\circ \underline{\underline{j}}$$

$$= \frac{3}{2} \underline{\underline{i}} + \frac{3\sqrt{3}}{2} \underline{\underline{j}} - \frac{4}{\sqrt{2}} \underline{\underline{i}} + \frac{4}{\sqrt{2}} \underline{\underline{j}}$$

$$= \left(\frac{3}{2} - \frac{4}{\sqrt{2}} \right) \underline{\underline{i}} + \left(\frac{3\sqrt{3}}{2} + \frac{4}{\sqrt{2}} \right) \underline{\underline{j}}$$

$$\begin{aligned} \therefore \underline{\underline{a}} &= \frac{\underline{\underline{\Sigma F}}}{m} = \frac{1}{2} \left(\frac{3}{2} - \frac{4}{\sqrt{2}} \right) \underline{\underline{i}} + \frac{1}{2} \left(\frac{3\sqrt{3}}{2} + \frac{4}{\sqrt{2}} \right) \underline{\underline{j}} \\ &\approx -0.66 \underline{\underline{i}} + 2.71 \underline{\underline{j}} \end{aligned}$$

Question 5



Local minima at 'a' and 'f'; local maximum at 'd'.

[A]

Points of inflexion at 'b' and 'e'.

[A]

Correct general shape.

[A]