

INSIGHT Trial Exam Paper

2010 SPECIALIST MATHEMATICS

Written examination 1

STI	ID.	EN	ו דו	NΔ	M	E.
	,,,,	רוגיו				٠,,

QUESTION AND ANSWER BOOK

Reading time: 15 minutes Writing time: 1 hour

Structure of book

Number of questions	Number of questions to be answered	Number of marks	
10	10	40	

- Students are permitted to bring the following items into the examination: pens, pencils, highlighters, erasers, sharpeners and rulers.
- Students are NOT permitted to bring sheets of paper, notes of any kind or white out liquid/tape into the examination.
- Calculators are not permitted in this examination.

Materials provided

- The question and answer book of 15 pages with a separate sheet of miscellaneous formulas.
- Working space is provided throughout this book.

Instructions

- Write your **name** in the box provided.
- Remove the formula sheet during reading time.
- You must answer the questions in English.

Students are NOT permitted to bring mobile phones or any other electronic devices into the examination.

This trial examination produced by Insight Publications is NOT an official VCAA paper for the 2010 Specialist Mathematics written examination 1.


This examination paper is licensed to be printed, photocopied or placed on the school intranet and used only within the confines of the purchasing school for examining their students. No trial examination or part thereof may be issued or passed on to any other party including other schools, practising or non-practising teachers, tutors, parents, websites or publishing agencies without the written consent of Insight Publications.

Copyright © Insight Publications 2010

The position of a particle is given by $r = \sec \frac{t}{2} \stackrel{i}{\sim} + 2 \tan \frac{t}{2} \stackrel{j}{\sim}$, where $t \in [0, \pi)$.


a. Show that the Cartesian equation of the particle is of the form $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. Find the value of a and b and state the domain and range.

b. Sketch the graph of its path on the axes below.



1 mark

a. On the set of axes below, shade the region enclosed by the graph of $y = \sin^{-1} x$ and the lines y = 0 and x = 1.



1 mark

b.	Find the exact area enclosed by the graph of $y = \sin^{-1} x$ and the lines $y = 0$	and $x = 1$.
		2 marks
		2 marks
c.	Find the exact volume generated by rotating this area about the <i>y</i> -axis.	
		2 1
		3 mark

_			_
On	est	io	n 3


Given that $z_1 = 2$ is a solution to the equation $z^2 + (i\sqrt{3} - 3)z + 2 - 2\sqrt{3}i = 0$, $z \in \mathbb{C}$ Show that the other solution is $z_2 = 1 - \sqrt{3} i$. 1 mark Find the square root of z_2 in the form a + bi. b.

$\boldsymbol{\cap}$		
	uestion	
ι,	ucsuvii	-

Find the equation of the tangent to the function $x \log_e y + 2x^2 = 3$ at the point where $x = 1$.				

Question	5
Question	J

£		
a.	Show that $\frac{2x}{4-x^2} = \frac{1}{2-x} - \frac{1}{2+x}$.	
b.	Hence, or otherwise, find $\int_{-1}^{1} \frac{2x}{4-x^2} dx$.	2 marks



Given that |a| = |b| in the trapezium shown above, use vectors to show that $\triangle ACD$ is right angled.


This page is blank

A parachutist free-falls from rest and reaches a speed of 2g m/s when his parachute is activated.

The acceleration of the parachutist from this point is (g - 2v) m/s², where v is the speed of the parachutist t seconds after the parachute is activated.

Five seconds after the parachute is activated, the speed of the parachutist is $\frac{g}{2}(a + be^{-10})$.				
Find the values of a and b . The acceleration due to gravity is $g = 9.8 \text{ m/s}^2$.				

A 2 kg mass on a horizontal bench is connected by an inextensible string via a smooth pulley to a 5 kg mass, as shown below. The force of *P* newtons is acting on the 2 kg mass, as shown. The coefficient of friction between the 2 kg mass and the bench is 0.25.



a.	Label	all	forces	on the	diagram	above
----	-------	-----	--------	--------	---------	-------

1 mark


b.	Show that the minimum value of the force P required to maintain equilibrium is $4.5g$ newtons.

If $P = 2.5g$, find the exact acceleration of the s	system.
	2 marks
Question 9	
Evaluate $\int_{e}^{e^2} \frac{1}{x \ln x} . dx.$	

S =	$\{z\cdot (z+$	$1)(\bar{z} +$	1) = 4	$z \in \mathbb{C}$	and	$T = \{z: Arg$	z(z+1)	$=\frac{-3\pi}{}$	$z \in$	C }
J	(2. (2)	1)(2	1) 7,	2 - 0;	ana	1 (2.711)	, (2 ' 1,	, , ,	<i>2</i> C	C 3 .

a.	Find the Cartesian equations of S and T .

b. Show the subsets S and T on the complex plane below.



1 mark